## **Appendix 4. Application References**

- AgResearch (2010). Managing parasite resistance to anthelmintic drugs. Science Review. http://www.agresearch.co.nz/publications/sciencereview/parasite-resistance.pdf. Retrieved 25 August, 2010.
- Bang HS, Lee J-H, Kwon OS, Na YE, Jang YS, Kim WH. (2005). Effects of paracoprid dung beetles (Coleoptera: Scarabaeidae) on the growth of pasture herbage and on the underlying soil. *Applied Soil Ecology* 29: 165–171.
- Bishop DM. (1998). Parasitic Hyrnenoptera reared from dung-breeding Diptera in New Zealand. *New Zealand Entomologist* 21: 99- 106.
- Bishop DM, Heath ACG. (1995). *Aphaereta aotea*, a hymenopteran parasitoid with the potential to act as a biological control agent agzinst flystrike. *New Zealand Journal of Zoology* 22: 187 (Abstract).
- Bishop DM, Heath ACG, Haack NA. (1996). Distribution, prevalence and host associations of hymenoptera parasitic on Calliphoridae occurring in flystrike in New Zealand. *Medical and Veterinary Entomology* 10: 365-370.
- Bornemissza GF. (1960). Could dung eating insects improve our pastures? *Journal of the Australian Institute of Agricultural Science* 26: 54-56.
- Bornemissza GF. (1970). Insectary studies on the control of dung breeding flies by the activity of the dung beetle, *Onthophagus gazella* F. (Coleoptera: Scarabaeinae). *Journal of the Australian Entomological Society* 9: 31-41.
- Bornemissza GF. (1976). The Australian dung beetle project 1965-1975. *Australian Meat Research Committee Review* 30: 32pp.
- Bornemissza GF, Williams CH. (1970). An effect of dung beetle activity on plant yield. *Pedobiologia* 10: 1-7.
- Blume RR, Matter JJ, Eschle JL. (1973). *Onthophagus gazella*: effect on survival of horn flies in the laboratory. *Environmental Entomology* 2: 811-813.
- Bryan RP. (1973). The effects of dung beetle activity on the numbers of parasitic gastrointestinal helminth larvae recovered from pasture samples. *Australian Journal of Agricultural Research* 24: 161-168.
- Bryan RP. (1976). The effect of the dung beetle, *Onthophagus gazella*, on the ecology of the infective larvae of gastrointestinal nematodes of cattle. *Australian Journal of Agricultural Research* 27: 567-574.
- Cambefort Y. (1991). From saprophagy to coprophagy. Pp. 22-35. In: Hanski I. & Cambefort Y. (Eds), *Dung Beetle Ecology*. Princeton University Press, Princeton, New Jersey.
- Cambefort Y, Hanski I. (1991). Dung beetle population biology. Pp. 36-50. In: Hanski I. & Cambefort Y. (Eds), *Dung Beetle Ecology*. Princeton University Press, Princeton, New Jersey.
- Cameron PJ, Butcher CF. (1979). Pitfall trapping and soil sampling for Scarabaeidae (Coleoptera) and possible predators in some Northland and Auckland pastures. Pp. 113-117. In: Crosby TK, Pottinger RP (Eds). Proceedings of the 2<sup>nd</sup> Australasian Conference on Grassland Invertebrate Ecology, Government Printer, Wellington.
- Cameron PJ, Thomas WP. (1989). *Heteronychus arator* (F.), black beetle (Coleoptera: Scarabaeidae). Pp. 17-21. In: Cameron PJ, Hill RJ, Bain J, Thomas WP (Eds) *A review of biological control of invertebrate pests and weeds in New Zealand 1874-1987*. Technical Bulletin No. 10, DSIR Entomology Division, Publ. by CAB International Wallingford, Oxford, UK.

Appendix 4 Page 1 of 6

- Cameron PJ, Wigley PJ. (1989). *Costelytra Zealandica* (White), grass grub (Coleoptera: Scarabaeidae). Pp. 9-16. In: Cameron PJ, Hill RJ, Bain J, Thomas WP (Eds) *A review of biological control of invertebrate pests and weeds in New Zealand 1874-1987*. Technical Bulletin No. 10, DSIR Entomology Division, Publ. by CAB International Wallingford, Oxford, UK.
- Chan KY, Barchia I. (2007). Soil compaction controls the abundance, biomass and distribution of earthworms in a single dairy farm in south-eastern Australia. *Soil and Tillage Research* 94: 75–82.
- Chirico J, Wiktelius S, Waller PJ (2003). Dung beetle activity and the development of trichostrongylid eggs into infective larvae in cattle faeces. *Veterinary Parasitology* 118(1-2): 157-163.
- Davis ALV. (1987). Geographical distribution of dung beetles (Coleoptera: Scarabaeidae) and their seasonal activity in south-western Cape Province. *Journal of the Entomological Society of Southern Africa* 50(2): 275-285.
- Davis ALV. (1996a). Seasonal dung beetle activity and dung dispersal in selected South African habitats: Implications for pasture improvement in Australia. *Agriculture Ecosystems & Environment* 58(2-3): 157-169.
- Davis ALV. (1996b). Community organization of dung beetles (Coleoptera: Scarabaeidae): Differences in body size and functional group structure between habitats. *African Journal of Ecology* 34(3): 258-275.
- Davis ALV, Scholtz CH. (2001). Historical versus ecological factors influencing global patterns of scarabaeine dung beetle diversity. *Diversity and Distributions* 7: 161-174.
- de Klein CAM, Sherlock RR, Cameron KC, van der Weerden TJ. (2001). Nitrous oxide emissions from agricultural soils in New Zealand—a review of current knowledge and directions for future research. *Journal of the Royal Society of New Zealand* 31: 543-574.
- Doube BM. (1983). The habitat preference of some bovine dung beetles (Coleoptera: Scarabaeidae) in Hluhluwe Game Reserve, South Africa. *Bulletin of Entomological Research* 73: 357-371.
- Doube BM. (1990). A functional classification for analysis of the structure of dung beetle assemblages. *Ecological Entomology* 15: 371-383.
- Doube BM. (2004). Dung beetle–Cryptosporidium research: review and a pilot study on the recovery of *C. parvum* oocysts from dung buried by the dung beetle *Bubas bison*. Prepared for Watershed Protection office, June: 29pp.
- Doube BM. (2005a). Optimal release sites for the dung beetle *Bubas bison* in the water catchments of the Central Highlands, Victoria. Report prepared for Central Highlands Water: 17pp.
- Doube BM. (2005b). Dung burial by the winter-active beetle *Bubas bison* and its impact on soil properties in the Barham River Water Catchment, Apollo Bay, Victoria. Final report for phase 1: 31pp.
- Doube BM. (2005c). The impact of the activity of the late summer/autumn-active dung beetle *Geotrupes spiniger* on soil and water properties in the field. Final Report. Dairy SA, February: 23pp.
- Doube BM. (2006a). The impact of dung burial by *Bubas bison* and *Geotrupes spiniger* on earthworm populations in the Central Highlands of Victoria. Final Report, Central Catchments Water: 23pp.
- Doube BM. (2006b). Adding value to a Dairy SA dung beetle project: Evaluation of pasture growth due to the late summer/autumn-active dung beetle *Geotrupes spiniger* at Flaxley SA. Final report, Dairy South Australia: 9pp.

Appendix 4 Page 2 of 6

- Doube BM. (2007). Pasture growth benefits of the dung beetle *Bubas bison* in the Margaret River region of Western Australia. Interim report for June 2007: Assessing pasture growth in the field trial at Margaret River. WA Water Corporation, September: 11pp.
- Doube BM. (2008a). The pasture growth and environmental benefits of dung beetles to the southern Australian cattle industry. Meat & Livestock Industry, North Sydney, NSW. ISBN: 9781741912814: 192pp.
- Doube BM. (2008b). Appendix3: Subsoil Chemical analysis, August and November 2006, May and September 2007. Pp. 71-104. In: *The pasture growth and environmental benefits of dung beetles to the southern Australian cattle industry*. Meat & Livestock Industry, North Sydney, NSW.
- Doube BM. (2008c). Appendix 5: The effect of the dung beetle I on earthworm abundance and distribution. Pp. 129-158. In: *The pasture growth and environmental benefits of dung beetles to the southern Australian cattle industry*. Meat & Livestock Industry, North Sydney, NSW.
- Duraes R, Martins WP, Vaz-De-Mello FZ (2005). Dung beetle (Coleoptera: Scarabaeidae) assemblages across a natural forest-cerrado ecotone in Minas Gerais, Brazil. *Neotropical Entomology* 34(5): 721-73.
- Dymock JJ. (1993). A case for the introduction of additional dung burying beetles (Coleoptera: Scarabaeidae) into New Zealand. *New Zealand Journal of Agricultural Research* 36: 163-171.
- Dymock JJ. (1989). *Listronotus bonariensis* (Kuschel), Argentine stem weevil (Coleoptera: Curculionidae). Pp. 23-26, In: Cameron PJ, Hill RJ, Bain J, Thomas WP (Eds) *A review of biological control of invertebrate pests and weeds in New Zealand 1874-1987*. Technical Bulletin No. 10. DSIR Entomology Division, Publ. by CAB International Wallingford, Oxford, UK.
- Edwards CA. (2004). The importance of earthworms as key representatives of the soil fauna. Pp 3-11. In: CA Edwards (Ed.) *Earthworm Ecology*, 2nd Edition. CRC Press, Boca Raton, Florida: 441pp.
- Edwards P. (2003). Improving sustainable land management systems in Queensland using Beetles: Final report of the 2001-2002 Queensland Dung Beetle Project: 55pp (pg 3).
- Edwards P. (2007). Introduced Dung Beetles in Australia 1967-2007: current status and future directions, Landcare Australia: 82pp.
- Edwards P. (2010). Biological Control of Pastoral Dung in New Zealand. A report on the climatic suitability of exotic dung beetle species for introduction to New Zealand: 39pp.
- Fay HAC, Macqueen A, Doube BM, Kerr JD. (1990). Impact of fauna on mortality and size of Haematobia spp. (Diptera: Muscidae) in dung pads in Australia and South Africa. *Bulletin of Entomological Research* 80: 385–392.
- Fincher GT. (1973). Dung beetles as biological control agents for gastrointestinal parasites of livestock. *Journal of Parasitology* 59 (2): 396-399.
- Fincher G T. (1975). Effects of dung beetle activity on the number of nematode parasites acquired by grazing cattle. *Journal of Parasitology* 61: 759-762.
- Fincher GT. (1981). The potential value of dung beetles in pasture ecosystems. *Journal of the Georgia Entomological Society* 16(1): 316-333.
- Fincher GT, Stewart TB. (1979). Vertical migration by nematode larvae of cattle parasites through soil. *Proceedings of the Helminthological Society*, Washington 46: 43-46.
- Fincher GT, Monson WG, Burton GW. (1981). Effects of cattle feces rapidly buried by dung beetles on yield and quality of coastal Bermuda grass. *Agronomy Journal* 73: 775-779.

Appendix 4 Page 3 of 6

- Forgie SA. (2009). Reproductive activity of *Onthophagus granulatus* Boheman (Coleoptera: Scarabaeinae) in New Zealand: Implications for its effectiveness in the control of pastoral dung. *New Zealand Entomologist* 32: 76-84.
- Funasaki GY, Lai P-Y, Nakahara LM, Beardsley JW, Ota AK. (1988). A Review of Biological Control Introductions in Hawaii: 1890 to 1985. *Proceedings, Hawaiian Entomological Society* 28: 105-160.
- Gillard P. (1967). Coprophagous beetles in pasture ecosystems. *The Journal of the Australian Institute of Agricultural Science*, March: 30-34.
- Hanski I. (1991). The dung insect community. Pp. 5-21. In: Hanski I. & Cambefort Y. (Eds), *Dung Beetle Ecology*. Princeton University Press, Princeton, New Jersey.
- Haynes RJ, Williams PH. (1993). Nutrient cycling and soil fertility in the grazed pasture ecosystem. *Advances in Agronomy* 49: 119-199.
- Holter P. (1997). Methane emissions from Danish cattle dung pats in the field. *Soil Biology and Biochemistry* 29(1): 31-37.
- Holter P, Scholtz CH, Wardaugh KG. (2002). Dung feeding in adult scarabaeines (tunnellers and endocoprids): even large dung beetles eat small particles. *Ecological Entomology* 27: 169-176.
- Howden HF, Nealis VG. (1975). Effects of clearing in tropical rain forest on the composition of the coprophagous scarab beetle fauna (Coleoptera). *Biotropica* 7(2): 77-83.
- Jarvis SC, Lovell RD, Panayides R. (1995). Patterns of methane emission from excreta of grazing animals. *Soil Biology and Biochemistry* 27(12): 1581-1588.
- Klein BC. (1989). Effects of forest fragmentation on dung and carrion beetle communities in central Amazonia. *Ecology* 70(6): 1715-1725.
- Krell F-T. (2000). The fossil record of Mesozoic and Tertiary Scarabaeoidea (Coleoptera: Polyphaga). *Invertebrate Taxonomy* 14, 871-905.
- Le Jambre LF. (2009). Dung beetles and internal parasites of sheep. Final Report prepared for Meat and Livestock Australia (SUPER PIRD ref.: S2005/NO3). Granite Borders Landcare Committee, Tenterfield, NSW, Australia: 21pp.
- Losey JE, Vaughan M. (2006). The Economic Value of Ecological Services Provided by Insects. *BioScience* 56(4), 311-323.
- MAF, no date. Livestock statistics. Ministry of Agriculture and Forestry. http://www.maf.govt.nz/statistics/pastoral/livestock-numbers/. Retrieved 26 August, 2010.
- Macqueen A, Beirne BP. (1975). Effects of cattle dung and dung beetle activity on growth of beardless wheatgrass in British Columbia. *Canadian Journal of Plant Science* 55: 961-967.
- Magura T, Tóthmérész B, Molnár T. (2001). Forest edge and diversity: carabids along forest-grassland transects. *Biodiversity and Conservation* 10: 287–300.
- Marten GC, Donker JD. (1966). Animal excrement as a factor influencing acceptability of grazed forage. In: Proceedings of the 10th International Grassland Congress, Helsinki: 359-363.
- Martin NA, Charles JC. (1979). Lumbricid earthworms and cattle dung in New Zealand pastures. In: Proceedings of the 2nd Australasian Conference on Grassland Invertebrate Ecology: 52-54.
- Mathison BA, Ditrich O. (1999). The fate of *Cryptosporidium parvum* oocysts ingested by dung beetles and their possible role in the dissemination of cryptosporidiosis. *Journal of Parasitology* 85: 678-681.

Appendix 4 Page 4 of 6

- MfE. (2010). Climate change information, New Zealand. Ministry for the Environment.

  <a href="http://www.climatechange.govt.nz/emissions-trading-scheme/participating/agriculture.html">http://www.climatechange.govt.nz/emissions-trading-scheme/participating/agriculture.html</a>
  Retrieved 26 August, 2010. No Article (pdf) available.
- Miller A. (1961). The mouthparts and digestive tract of adult dung beetles (Coleoptera: Scarabeidae) with reference to the ingestion of helminth eggs. *Journal of Parasitology* 47:735–744.
- Miller A, Chi-Rodriquez E, Nichols RL. (1961). The fate of helminth eggs and protozoan cysts in human feces ingested by dung beetles (Coleoptera: Scarabeidae). *American Journal of Tropical Medicine and Hygiene* 10: 748–754.
- Nelson W, Harris B. (2006). Flies, fingers, fomites, and food. Campylobacteriosis in New Zealand\_food-associated rather than food-borne. *The New Zealand Medical Journal* 119 (1240): 1-7.
- Nichols E, Spector S, Louzada J, Larsen T, Amezquita S, Favila ME. (2008). Ecological functions and ecosystem services provided by Scarabaeinae dung beetles. *Biological Conservation* 141: 1461-1474.
- Norman R. (2010). Time for Governments to regulate dairy pollution. Press Release. Green Party of Aotearoa New Zealand. http://www.greens.org.nz/press-releases/time-government-regulate-dairy-pollution. Retrieved 24 August, 2010.
- Osberg DC, Hanrahan SA, Doube BM. (1992). The spatial distribution of *Allogymnopleurus thalassinus* Klug and *A. consocius* Peringuey (Coleoptera: Scarabaeidae) in an area of mixed soil types in South Africa. *Journal of the Entomological Society of Southern Africa* 55(1): 85-92.
- Pawson SM, Brockerhoff EG, Meenken ED, Didham RK. (2008). Non-native plantation forests as alternative habitat for native forest beetles in a heavily modified landscape. *Biodiversity and Conservation* 17:1127–1148.
- Popay A, Marshall S. (1996). The effect of the dung beetles Onthophagus posticus and Onthophagus granulatus on survival of parasitic nematodes in sheep dung. AgResearch report for MRDC Project, number 96PR 36/1.1: 5pp.
- Rattray PV. (2003). Internal parasite report. Meat and Wool Innovation Ltd, contract to Merino New Zealand Inc, Christchurch. October: 187pp.
- Richardson P, Richardson R. (2000). Dung beetles and their effects on soil. *Ecological Restoration* **18**:116-117. URL: managingwholes.com/dung-beetles.htm. Site last visited: 28July, 2010.
- Ridsdill-Smith TJ, Edwards PB. Biological control: ecosystem functions provided by dung beetles Chapter 12. In: Simmons LW, Ridsdill-Smith TJ (Eds), *Ecology and evolution of dung beetles*. Wiley Blackwell due to be published in April 2011. *Chapter currently not available*.
- Ridsdill-Smith TJ, Matthiessen JN. (1988). Bush fly, Musca vetustissima Walker (Diptera, Muscidae), control in relation to seasonal abundance of scarabaeinae dung beetle affix (Coleoptera: Scarabaeidae) in south-western Australia. *Bulletin of Entomological Research* 78: 633-639.
- Römbke J, Jänsch S, Didden W. (2005). The use of earthworms in ecological soil classification and assessment concepts. *Ecotoxicology and Environmental Safety* 62: 249–265.
- Rougon D, Rougon C, Levieux J, Trichet J. (1990). Variations in the amino-acid content in zebu dung in the Sahel during nesting by dung beetles (Coleoptera: Scarabaeidae). *Soil Biology and Biochemistry* 22(2): 717-723.
- Scholtz CH, Chown SL. (1995). The evolution of habitat use and diet in the Scarabaeoidea: A phylogenetic approach. Pp. 354-374. In: Pakaluk J, Slipinski SA (Eds), *Biology, Phylogeny and Classification of Coleoptera: Papers celebrating the 80th birthday of Roy A. Crowson*. Museum I Instytut Zoologii PAN, Warszawa.

Appendix 4 Page 5 of 6

- Seldon DS. (2002). A comparison of ground insect distributions from several New Zealand fragmented forest habitats. BSc(Hons) Dissertation, University of Auckland: 60pp.
- Spector S, Ayzama S (2003). Rapid turnover and edge effects in dung beetle assemblages (Scarabaeidae) at a Bolivian Neotropical forest-savanna ecotone. *Biotropica* 35(3): 394-404.
- Strong L. (1992). Avermectins: A review of their impact on insects of cattle dung. *Bulletin of Entomological Research* 82: 265-274.
- Stufkens MW, Farrell JA. (1989). *Sitona discoideus* Gyllenhal, sitona weevil (Coleoptera: Curculionidae). Pp. 27-28. In: Cameron PJ, Hill RJ, Bain J, Thomas WP (Eds) *A review of biological control of invertebrate pests and weeds in New Zealand 1874-1987*. Technical Bulletin No. 10, DSIR Entomology Division, Publ. by CAB International Wallingford, Oxford, UK.
- Waghorn TS, Leathwick DM, Chen L-Y, Gray RAJ, Skipp RA. (2002). Influence of nematophagous fungi, earthworms and dung burial on development of the free-living stages of *Ostertagia* (*Teladorsagia*) *circumcincta* in New Zealand. *Veterinary Parasitology* 104: 119–129.
- Wardhaugh K.G. (2005). Insecticidal activity of synthetic pyrethroids, organophosphates, insect growth regulators, and other livestock parasiticides: an Australian perspective. *Environmental Toxicology and Chemistry* 24 (4): 789-796.
- Waterhouse DF. (1974). The biological control of dung. Scientific American 230: 100-109.
- Whitehead D. (2009). A dairy farm is not greenhouse gas neutral. Landcare Research article prepared for Dairy NZ Ltd, <a href="http://www.carbonzero.co.nz/publications/dairyNZ">http://www.carbonzero.co.nz/publications/dairyNZ</a> mar09.pdf: 4 pp.
- Williams PH, Haynes RJ. (1995). Effect of sheep, deer and cattle dung on herbage production and soil nutrient content. *Grass and Forage Science* 50: 263-271.
- Yokoyama K, Kai H, Tsuchiyama H. (1991). Paracoprid dung beetles and gaseous loss of nitrogen from cow dung. *Soil Biology and Biochemistry* 23: 643–647.

Appendix 4 Page 6 of 6